

# A knot Floer stable homotopy type

Ciprian Manolescu  
*based on joint work with Sucharit Sarkar*

February 1, 2022

To a grid diagram  $\mathbb{G}$ , we will associate a stable homotopy type whose homology is grid homology (link Floer homology).

- 1 Review of grid homology
- 2 Statement of results
- 3 Framed flow categories
- 4 The inductive construction

**Ozsváth-Szabó, Rasmussen**, 2003: *knot Floer homology*  $HFK =$  bigraded homology theory for knots. Different versions:  $\widehat{HFK}$ ,  $HFK^-$ ,  $HFK^+$ , etc.

$\sum_{i,j} (-1)^i q^j \widehat{HFK}_i(K, j) = \Delta_K(q)$ , the Alexander polynomial

**Generalization:**  $HFL = link Floer homology$  (**Ozsváth-Szabó**, 2005)

It detects fiberedness, Thurston norm, has many applications to concordance, surgery questions, etc.


**Combinatorial description:** *grid homology* (**M.-Ozsváth-Sarkar**, 2006), in terms of grid diagrams.

Different versions:  $\widehat{GH}$ ,  $GH^+$ ,  $GH^-$ , etc. We will focus on  $GH^+$ .


# Grid diagrams

Every link in  $S^3$  admits a *grid diagram*  $\mathbb{G}$ ; that is, an  $n$ -by- $n$  grid on the torus with  $O$  and  $X$  markings inside such that:

- Each row and each column contains exactly one  $X$  and one  $O$ ;
- As we trace the vertical and horizontal segments between  $O$ 's and  $X$ 's (verticals on top), we see the link  $L$ .




# States




We define a state  $x = \{x_1, \dots, x_n\}$  to be an  $n$ -tuple of points on the grid (one on each vertical and horizontal circle). The set of states is denoted  $\mathbb{S}$ .

# Generators



We define a state  $x = \{x_1, \dots, x_n\}$  to be an  $n$ -tuple of points on the grid (one on each vertical and horizontal circle). The set of states is denoted  $\mathbb{S}$ .

# Differentials



In a grid diagram  $\mathbb{G}$ , index 1 pseudo-holomorphic strips in  $\text{Sym}^n(\mathbb{G})$  are in 1-to-1 correspondence to empty rectangles on the grid (that is, having no red or blue dots inside).

# The grid complex

Different ways of keeping track of the  $O$  and  $X$  markings  $\rightsquigarrow$  different versions of the grid complex.

The version  $GC^+(\mathbb{G})$  is generated by

$$U_1^{-j_1} \dots U_n^{-j_n} x = [x, j_1, \dots, j_n], \quad x \in \mathbb{S}, \quad j_1, \dots, j_n \in \mathbb{N}$$

in homological grading  $\text{gr}(x) + 2j_1 + \dots + 2j_n$ .

The differential on  $GC^+$  is given by

$$\partial([x, j_1, \dots, j_n]) = \sum_y \sum_{\substack{R \in \mathcal{R}(x, y) \\ \mathbb{X}(R) = (0, \dots, 0)}} s(R) U^{\mathbb{O}(R)} [y, j_1, \dots, j_n],$$

where  $U^{\mathbb{O}(R)} := U_1^{O_1(R)} \dots U_n^{O_n(R)}$ .

The homology  $GH^+(\mathbb{G}) = GH^+(K) = HFL^+(K)$  is independent of  $\mathbb{G}$ .



Grid homology is an example of **Lagrangian Floer homology**:

$M = \text{Sym}^n(\mathbb{G})$  symplectic manifold;  $L_0 = \mathbb{T}_\alpha$ ,  $L_1 = \mathbb{T}_\beta$  Lagrangians

$\rightsquigarrow \mathcal{M}(x, y)$  moduli spaces of pseudo-holomorphic strips

$\rightsquigarrow (CF(L_0, L_1), \partial) \rightsquigarrow HF(L_0, L_1)$ . To have  $\partial^2 = 0$ , ideally we want

$$\partial \overline{\mathcal{M}}(x, y) = \coprod_z \overline{\mathcal{M}}(x, z) \times \overline{\mathcal{M}}(z, y).$$

In general, in symplectic geometry, the compactification  $\overline{\mathcal{M}}(x, y)$  has additional points, coming from **disk and sphere bubbles**. Thus, we may have  $\partial^2 \neq 0$ .

In the situation at hand ( $GH^+$ ), we do not have sphere bubbles, but we do have disk bubbles, corresponding to rows and columns on the grid. We still have  $\partial^2 = 0$ , because the row through  $O_i$  cancels out the column through  $O_j$ .

# The Spanier-Whitehead suspension category

*Objects* (suspension spectra): pairs  $(X, n) = \Sigma^{-n}X$ , where  $X =$  pointed CW complex,  $n \in \mathbb{Z}$ .

*Morphisms* are stable homotopy classes of maps:

$$\text{Hom}((X, n), (Y, m)) = \text{colim}_{q \rightarrow \infty} [\Sigma^{q-n}X, \Sigma^{q-m}Y].$$

e.g.  $(S^0, 5) = S^{-5}$  the “ $(-5)$ -dimensional sphere”

The (reduced) homology of  $(X, n)$  is  $\tilde{H}_i(X, n) = \tilde{H}_{i+n}(X)$ .

**Generalizations:** spectra, pro-spectra, etc.

# A knot Floer stable homotopy type

To each grid diagram  $\mathbb{G}$  and integer  $j$  we will associate a suspension spectrum  $\mathcal{X}_j^+(\mathbb{G})$  such that

$$\tilde{H}_i(\mathcal{X}_j^+(\mathbb{G}); \mathbb{Z}) = GH_{i,j}^+(\mathbb{G}).$$


Just as  $GH^+$  comes equipped with the structure of a module over the polynomial ring  $\mathbb{Z}[U_1, \dots, U_n]$ , here we have maps

$$U_i : \mathcal{X}_j^+(\mathbb{G}) \rightarrow \Sigma^2 \mathcal{X}_{j-1}^+(\mathbb{G}),$$

where  $\Sigma^2$  denotes the double suspension.

The construction is based on *framed flow categories* (**Cohen-Jones-Segal**, 1995) and similar in spirit to the construction of *Khovanov stable homotopy types* by **Lipshitz-Sarkar** (2011).

# An example: $T(2, 5)$



$CFK^+(T(2, 5))$

In gradings  $A = j \neq 1$ , the grid homology  $GH_j^+$  is supported in at most two consecutive gradings  $\Rightarrow \mathcal{X}_j^+$  is a wedge of spheres; e.g.  $\mathcal{X}_{-1}^+ = S^{-3} \vee S^{-2}$ .

When  $j = 1$ , the spectrum  $\mathcal{X}_{-1}^+$  has two cells in dimensions  $-1$  and  $2$ , attached via a map  $\tau : S^1 \rightarrow S^{-1}$ . There are two possibilities, according to  $[\tau] \in \pi_2^{st}(S^0) = \mathbb{Z}/2$ .

We expect that  $[\tau] = 0$ , and therefore  $\mathcal{X}_1^+ = S^{-1} \vee S^2$ .

We can do similar constructions for other versions of grid homology ( $\widehat{GH}$ ,  $GH^-$ , ...) provided we don't allow the domains of pseudo-holomorphic strips to cross a particular  $X$ -marking on the grid.

## Still to do:

- Prove that the stable homotopy type of  $\mathcal{X}_j^+(\mathbb{G})$  is a knot invariant;
- Construct versions going over all  $O$  and  $X$ -markings;
- Constructions using pseudo-holomorphic curves ( $\rightsquigarrow$  Heegaard Floer stable homotopy types);
- Non-trivial computations: examples where  $GH_j^+(K) = GH_j^+(K')$  but  $\mathcal{X}_j^+(K) \neq \mathcal{X}_j^+(K')$ .

# Framed flow categories

To upgrade a chain complex  $C_*$  to a suspension spectrum, we need a *framed flow category* (cf. **Cohen-Jones-Segal**):

**Objects** = generators of  $C_*$

**Morphisms:**  $\text{Hom}(x, y) = \overline{\mathcal{M}}(x, y)$  a compact manifold with corners (in fact,  $\langle d \rangle$ -manifold) of dimension  $d = \text{gr}(x) - \text{gr}(y) - 1$ ; e.g. in Lagrangian  $HF$ , the compactified moduli space of holomorphic strips from  $x$  to  $y$ .

We require that

$$\partial \overline{\mathcal{M}}(x, y) = \coprod_z \overline{\mathcal{M}}(x, z) \times \overline{\mathcal{M}}(z, y)$$

- + neat embeddings of  $\overline{\mathcal{M}}(x, y)$  into suitable  $\mathbb{R}_+^k \times \mathbb{R}^m$
- + framings of their normal bundles
- + compatibility conditions.

# From framed flow categories to spectra

From a framed flow category we get a spectrum  $\mathcal{X}$  by attaching one  $d$ -dimensional cell for each generator in degree  $d$ .

The spaces  $\overline{\mathcal{M}}(x, y)$  tell us the attaching maps, via the Pontryagin-Thom construction.

**Example:** Say we have two generators  $y$  and  $x$  in degrees  $a < b$ . Let  $k = b - a - 1$ . Then  $\mathcal{X}$  is obtained from  $S^a$  by attaching a  $b$ -cell via a stable map

$$\tau : S^{b-1} \rightarrow S^a, \quad [\tau] \in \pi_k^{st}(S^0) \cong \Omega_{fr}^k$$

where  $\Omega_{fr}^k$  is the framed cobordism group.

The stably framed manifold  $\overline{\mathcal{M}}(x, y)$  gives the desired element in  $\Omega_{fr}^k$ .

# A framed flow category for $GC_j^+$

To do this for the grid complex  $GC_j^+$ , we need to construct manifolds with corners

$$\overline{\mathcal{M}}(U_1^{i_1} \dots U_n^{i_n} x, U_1^{j_1} \dots U_n^{j_n} y) = \overline{\mathcal{M}}(x, U_1^{j_1 - i_1} \dots U_n^{j_n - i_n} y)$$

We do this whenever there is a positive domain (sum of rectangles)  $D$  from  $x$  to  $y$  going over the  $O$  basepoints a specified number of times.

Two domains are *equivalent* if they differ by a periodic domain, i.e. a linear combination of (row – column going through the same  $O_j$ ).

We write  $\overline{\mathcal{M}}(x, U_1^{j_1} \dots U_n^{j_n} y) = \overline{\mathcal{M}}([D])$ , where  $[D]$  is an equivalence class of positive domains.

This is a model for the moduli space of pseudo-holomorphic strips in  $\text{Sym}^n(\mathbb{C})$  supported in domains in the class  $[D]$ .



# Structure of the moduli spaces

Fix generators  $x$  and  $U_1^{j_1} \dots U_n^{j_n} y$ ; let  $D_1, D_2, \dots, D_k$  be all the positive domains that connect them (not passing through the special  $X$  marking and passing through  $O_i$   $j_i$  times), and let  $[D]$  be their equivalence class. To apply Cohen-Jones-Segal construction, we need the following:

- $d$ -dimensional  $\langle d \rangle$ -manifold  $\overline{\mathcal{M}}([D])$  (where  $d = \mu([D]) - 1$ ), with  $\partial_i \overline{\mathcal{M}}([D])$  (the  $i$ -th portion of the boundary) identified with products  $\overline{\mathcal{M}}([E]) \times \overline{\mathcal{M}}([F])$  with  $[E] + [F] = [D]$  and  $\mu([E]) = i$ .
- Neat embeddings of these moduli spaces into suitable Euclidean spaces, and coherent framings of their normal bundles.

# Gluing smaller pieces of the moduli spaces



What we do is construct some moduli spaces of the individual domains,  $\overline{\mathcal{M}}(D_i)$ . We then glue them together to get the space we want,

$$\overline{\mathcal{M}}([D]) = \overline{\mathcal{M}}(D_1) \cup \overline{\mathcal{M}}(D_2) \cup \cdots \cup \overline{\mathcal{M}}(D_k).$$


There are several issues:

- We neither expect nor construct these individual pieces  $\overline{\mathcal{M}}(D_i)$  as manifolds with corners; rather they are Whitney stratified spaces, with the stratification coming from disk bubbling.
- In order to glue these pieces, we need to keep track of these stratifications; consequently we need to keep track of a larger collection of moduli spaces—moduli spaces of domains along with a collection of disk bubbles attached to them. These will be the spaces  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$  from later in the talk.



# Example 1




# Example 2





# Example 3





# Example 4



# Example 5 (Part 1 of 4)





# Example 6 (Part 1 of 3)





# Example 6 (Part 2 of 3)




# The Whitney umbrella

In Example 6, the gluing is locally modeled on the decomposition of  $\mathbb{R}^3$  coming from the Whitney umbrella:

$$Z(2, 0, 0) = \{(a, b, c) \in \mathbb{R}^3 \mid a^2 b > c^2, c < 0\},$$

$$Z(1, 0, 1) = \{(a, b, c) \in \mathbb{R}^3 \mid a^2 b < c^2\},$$

$$Z(0, 0, 2) = \{(a, b, c) \in \mathbb{R}^3 \mid a^2 b > c^2, c > 0\},$$



Consider

$$\mathrm{Sym}^N(\mathbb{C})/\mathbb{R} \cong \mathbb{C}^N/\mathbb{R} \cong \mathbb{R}^{2N-1}$$

This decomposes according to how many of the imaginary parts of

$$\{z_1, \dots, z_N\} \in \mathrm{Sym}^N(\mathbb{C})$$

are negative or positive (or zero).

When  $N = 2$ , we recover the Whitney umbrella decomposition of  $\mathbb{R}^3$ .

More generally, we will encounter these decompositions of  $\mathbb{R}^{2N-1}$  as local models for our stratified spaces.

# Construction of the moduli spaces

We will construct moduli spaces

$$\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$$

which are models for the moduli spaces of pseudo-holomorphic strips with domain  $D$  and disk bubbles attached according to vectors


$$\vec{N} = (N_1, \dots, N_n), \quad \vec{\lambda} = (\lambda_1, \dots, \lambda_n)$$

Here  $N_j \in \mathbb{N}$ , and  $\lambda_j$  is an ordered partition of  $N_j$ . The number  $N_j$  counts the bubbles going through the  $j$ th  $O$ -marking. These bubbles are grouped according to the partition  $\lambda_j$ , with those in the same part appearing at the same height on the boundary of the pseudo-holomorphic strip.

The spaces  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$  will have specified local models, and come equipped with *neat embeddings* in  $\mathbb{R}_+^k \times \mathbb{R}^m$  and with normal framings.

# Permutohedra

We first construct  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$  when  $D = c_x$  (the trivial domain from some fixed  $x \in \mathbb{S}$  to itself), and all the entries of  $\vec{N}$  are 0's and 1's. In this case we define  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$  to be the permutohedron  $P_n$ , and give it a normal framing:



# The inductive argument

We define the rest of the spaces  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$  inductively on their dimension  $k$ . For the base case  $k = 0$ , we define them to be points.

For the inductive step, we suppose all spaces up to dimension  $k$  have been constructed, along with their embeddings and normal framings.

To construct a  $(k + 1)$ -dimensional space  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$ , we start with its (already constructed) boundary  $\partial \overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$  and smooth it to get a  $k$ -dimensional framed manifold  $\partial' \overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$ .

From here we get an element in the framed cobordism group

$$[\partial' \overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)] \in \Omega_{\text{fr}}^k.$$

# The complex of positive domains with partitions

We define a complex  $CDP$  whose generators are “positive domains with partitions,” i.e., triples  $(D, \vec{N}, \vec{\lambda})$ , where  $D$  does not go over a specified  $X$ -marking. The differential  $\delta : CDP_k \rightarrow CDP_{k-1}$  has four kinds of terms, corresponding to different boundaries of  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$ :

- **Type I** terms, given by subtracting a rectangle from  $D$ ;
- **Type II** terms, given by boundary degenerations, i.e., subtracting the row or the column through  $O_j$  from  $D$ , and at the same time increasing  $N_j$  by one, and changing  $\lambda_j$  accordingly; e.g.  $10 = 2 + 5 + 3 \rightarrow 11 = 2 + 5 + 1 + 3$ .
- **Type III** terms, given by combining two terms in one of the partitions  $\lambda_j$ ; e.g.  $10 = 2 + 5 + 3 \rightarrow 10 = 2 + 8$ . This corresponds to two bubbles reaching the same height.
- **Type IV** terms, given by dropping the first or final term in one of the partitions  $\lambda_j$ ; e.g.  $10 = 2 + 5 + 3 \rightarrow 7 = 2 + 5$ . This corresponds to removing a boundary degeneration, in the limit as its height goes to  $-\infty$  or  $+\infty$ .

# The complex of positive domains with partitions

## Proposition

For a grid diagram  $\mathbb{G}$  of size  $n$ , the homology of  $CDP_*$  is isomorphic to  $\mathbb{Z}^{2^n}$ . Its rank in degree  $k$  is  $\binom{n}{k}$ .

The homology is supported by the subcomplex  $CDP_*^\dagger \subset CDP_*$  the subcomplex generated by triples  $(c_x, \vec{N}, \vec{\lambda})$  where  $\vec{N}$  made only of 0's and 1's.

Let  $CDP'_* = CDP_*/CDP_*^\dagger$ .

## Proposition

The complex  $CDP'$  is acyclic (has trivial homology).



## Back to the construction

Altogether, the classes  $[\partial' \overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)] \in \Omega_{\text{fr}}^k$  produce an obstruction class

$$\sigma_k \in \text{Hom}(CDP'_{k+1}, \Omega_{\text{fr}}^k), \quad \sigma_k(D, \vec{N}, \vec{\lambda}) = [\partial' \overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)].$$

### Proposition

*The class  $\sigma_k$  is a cocycle.*

Since  $CDP'$  is acyclic, so is  $\text{Hom}(CDP'_{k+1}, \Omega_{\text{fr}}^k)$ . It follows that  $\sigma_k$  is the coboundary of some element  $\mathfrak{b} \in \text{Hom}(CDP'_k, \Omega_{\text{fr}}^k)$ .

We use  $\mathfrak{b}$  to adjust the definition of the  $k$ -dimensional moduli spaces that we previously constructed, so that all cocycles  $\sigma_k$  vanish. For this, we simply take disjoint unions with framed manifolds representing  $-\mathfrak{b}$ . (Note that do not change the definition of any moduli spaces of dimension  $k-1$  or lower.)

## Finishing the inductive step

Then  $\partial' \overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$  is framed null-cobordant. We fill it in arbitrarily to obtain the desired framed moduli space  $\overline{\mathcal{M}}_{\vec{N}, \vec{\lambda}}(D)$ , and continue with the induction.

**Observation:** It was essential that the complex  $CDP'$  be acyclic.

To arrange this:

- We had to avoid one  $X$ -marking;
- We had to first construct the moduli spaces for triples  $(D, \vec{N}, \vec{\lambda})$  where  $D = c_x$  and  $\vec{N}$  is made of 0's and 1's;
- We had to consider bubble configurations. Note that  $GC^+$  only involves domains that do not cross any  $X$ -marking, and therefore cannot contain a full row or column (so no bubbles). However, the analogue of  $CDP'$  where we avoid all  $X$ -markings has complicated homology.